

Quantum Grothendieck ring isomorphisms for quantum affine algebras of type A and B

Hironori OYA

Université Paris Diderot, IMJ-PRG

Based on a joint work with David HERNANDEZ

Oberseminar Algebra
Universität zu Köln, June 26, 2018

Motivation (1)

Topic : Finite dimensional representations of affine quantum groups

Question 1

Dimensions/ q -characters of simple modules ?

- \exists Classification of simple modules [Chari-Pressley 1990's]
“Highest weight theory”
- However, there are NO known closed formulae of their dimensions and q -characters in general. (e.g. \mathbb{A} analogue of Weyl-Kac character formulae...)

Question 2

Description of representation rings and their “deformations” ?

- Some (deformed) representation rings are known to be described nicely as (quantum) cluster algebras...

Motivation (2)

Question 1

Dimensions/ q -characters of simple modules ?

- ADE case \exists algorithm to compute them ! [Nakajima '04]
“Kazhdan-Lusztig algorithm”

The tool is t -deformed q -character, and the geometric construction (via quiver varieties) of simple modules guarantees this algorithm.

- Arbitrary (untwisted) case [Hernandez '04]
 - \exists t -deformed q -characters, **defined algebraically**
(\mathcal{A} geometry for non-symmetric cases)
 - Kazhdan-Lusztig algorithm gives **conjectural** q -characters of simple modules

However, they are still **candidates** in non-symmetric cases.

Motivation (3)

Question 2

Description of representation rings and their “deformations” ?

- [Hernandez-Leclerc '10 –, Kang-Kashiwara-Kim-Oh '15, Oh-Suh '16] The category of finite dimensional modules of affine quantum groups has several interesting monoidal subcategories ($\mathcal{C}_{\mathbb{Z}}$, $\mathcal{C}_{\mathbb{Z}}^{-}$, \mathcal{C}_{ℓ} , $\ell \in \mathbb{Z}$, $\mathcal{C}_{\mathcal{Q}}$ etc.), which are expected to be “monoidal categorifications” of cluster algebras (this fact is indeed proved in many cases).

Motivation (3)

Question 2

Description of representation rings and their “deformations” ?

- $X = \text{ADE}$ case Let
 - $K_t(\mathcal{C}_{\mathcal{Q}, X_n^{(1)}})$ the t -deformed Grothendieck ring (=quantum Grothendieck ring) of $\mathcal{C}_{\mathcal{Q}, X_n^{(1)}}$ for type $X_n^{(1)}$
 - $\mathcal{A}_v[N_-^{X_n}]$ the quantized coordinate algebra of the unipotent group of type X_n (\exists quantum cluster algebra structure !)(Each terminology will be explained later.)

Theorem (Hernandez-Leclerc '15)

$$K_t(\mathcal{C}_{\mathcal{Q}, X_n^{(1)}}) \simeq \mathcal{A}_v[N_-^{X_n}], \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} \leftrightarrow \text{dual canonical basis.}$$

Does it also hold in non-symmetric cases ?

Overview of Main results

In this talk, we consider the case of type $B_n^{(1)}$. Let $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ be the monoidal subcategory $\mathcal{C}_{\mathcal{Q}}$ for type $B_n^{(1)}$.

Theorem (Hernandez-O.)

$$\begin{array}{ccc} K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) & \simeq & \mathcal{A}_v[N_-^{A_{2n-1}}] & \stackrel{[\text{HL}]}{\simeq} & K_t(\mathcal{C}_{\mathcal{Q}', A_{2n-1}^{(1)}}) \\ \cup & & \cup & & \cup \\ \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} & \leftrightarrow & \text{dual canonical basis} & \stackrel{[\text{HL}]}{\longleftrightarrow} & \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} \end{array}$$

Remark

There are no known direct relations between the quantum affine algebras of type $B_n^{(1)}$ and $A_{2n-1}^{(1)}$ themselves.

Overview of Main results

In this talk, we consider the case of type $B_n^{(1)}$. Let $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ be the monoidal subcategory $\mathcal{C}_{\mathcal{Q}}$ for type $B_n^{(1)}$.

Theorem (Hernandez-O.)

$$\begin{array}{ccc} K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) & \simeq & \mathcal{A}_v[N_-^{A_{2n-1}}] & \stackrel{[\text{HL}]}{\simeq} & K_t(\mathcal{C}_{\mathcal{Q}', A_{2n-1}^{(1)}}) \\ \cup & & \cup & & \cup \\ \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} & \leftrightarrow & \text{dual canonical basis} & \stackrel{[\text{HL}]}{\longleftrightarrow} & \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} \end{array}$$

Kashiwara-Oh established an isomorphism between $K_{t=1}(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$ and $\mathbb{C}[N_-^{A_{2n-1}}]$ by a different method. Combining this result with our theorem above, we obtain the following :

Overview of Main results

Let $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ be the monoidal subcategory $\mathcal{C}_{\mathcal{Q}}$ for type $B_n^{(1)}$.

Theorem (Hernandez-O.)

$$\begin{array}{ccc}
 K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) & \simeq & \mathcal{A}_v[N_-^{A_{2n-1}}] & \xrightarrow{[\text{HL}]} & K_t(\mathcal{C}_{\mathcal{Q}', A_{2n-1}^{(1)}}) \\
 \cup & & \cup & & \cup \\
 \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} & \leftrightarrow & \text{dual canonical basis} & \xleftrightarrow{[\text{HL}]} & \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\}
 \end{array}$$

Kashiwara-Oh established an isomorphism between $K_{t=1}(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$ and $\mathbb{C}[N_-^{A_{2n-1}}]$ by a different method. Combining this result with our theorem above, we obtain the following :

Theorem (Hernandez-O.)

The (q, t) -characters of simple modules in $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ specialize to the corresponding q -characters.

Overview of Main results

Let $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ be the monoidal subcategory $\mathcal{C}_{\mathcal{Q}}$ for type $B_n^{(1)}$.

Theorem (Hernandez-O.)

$$\begin{array}{ccc} K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) & \simeq & \mathcal{A}_v[N_-^{A_{2n-1}}] & \stackrel{[\text{HL}]}{\simeq} & K_t(\mathcal{C}_{\mathcal{Q}', A_{2n-1}^{(1)}}) \\ \cup & & \cup & & \cup \\ \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} & \leftrightarrow & \text{dual canonical basis} & \stackrel{[\text{HL}]}{\longleftrightarrow} & \left\{ \begin{array}{l} (q, t)\text{-characters of} \\ \text{simple modules} \end{array} \right\} \end{array}$$

Theorem (Hernandez-O.)

The (q, t) -characters of simple modules in $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ specialize to the corresponding q -characters.

\rightsquigarrow The Kazhdan-Lusztig algorithm gives “correct” answers in $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$!

Quantum affine algebras

Let

- \mathfrak{g} a finite dimensional simple Lie algebra / \mathbb{C}
- $\mathcal{L}\mathfrak{g} := \mathfrak{g} \otimes_{\mathbb{C}} \mathbb{C}[t^{\pm 1}]$ its loop algebra $[X \otimes t^m, Y \otimes t^{m'}] = [X, Y] \otimes t^{m+m'}$
- $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ the Drinfeld-Jimbo quantum loop algebra / \mathbb{C} with a parameter $q \in \mathbb{C}^{\times}$ not a root of unity
generators : $\{k_i^{\pm 1}, x_{i,r}^{\pm}, h_{i,s} \mid i \in I, r \in \mathbb{Z}, s \in \mathbb{Z} \setminus \{0\}\}$

Properties

- $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ has a Hopf algebra structure.
- $\mathcal{U}_q(\mathfrak{g}) \xrightarrow[\text{Hopf alg.}]{} \mathcal{U}_q(\mathcal{L}\mathfrak{g}), e_i \mapsto x_{i,0}^+, f_i \mapsto x_{i,0}^-, k_i^{\pm 1} \mapsto k_i^{\pm 1}.$

Let \mathcal{C} be the category of finite-dimensional $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ -modules of type 1 (i.e. the eigenvalues of the actions of $\{k_i \mid i \in I\}$ are of the form $q^m, m \in \mathbb{Z}$).

Remark : \mathcal{C} is a non-semisimple abelian \otimes -category.

q -characters (1)

Let $V \in \mathcal{C}$. Frenkel-Reshetikhin showed that

$$\left\{ \begin{array}{l} \text{Generalized simultaneous eigenvalues of all } k_i^{\pm 1}, h_{i,s} \curvearrowright V \\ \text{Laurent monomials } m \text{ in } Y_{i,a} \text{'s } (i \in I, a \in \mathbb{C}^\times) \end{array} \right\} \overset{1:1}{\longleftrightarrow}$$

$\rightsquigarrow V = \bigoplus_m V_m$, called the ℓ -weight space decomposition.

$Y_{i,a}$ is an “affine analogue” of e^{ϖ_i} , ϖ_i fundamental weight.

Define the q -character of V as

$$\chi_q(V) := \sum_m \dim(V_m) m.$$

Then χ_q defines an injective algebra homomorphism

$$\chi_q: K(\mathcal{C}) \rightarrow \mathbb{Z}[Y_{i,a}^{\pm 1} \mid i \in I, a \in \mathbb{C}^\times] =: \mathcal{Y}_{\mathcal{C}^\times},$$

here $K(\mathcal{C})$ be the Grothendieck ring of \mathcal{C} [Frenkel-Reshetikhin].

$K(\mathcal{C})$ is commutative. (However sometimes $V \otimes W \not\cong W \otimes V$ in \mathcal{C} .)

q -characters (2)

Set $\mathcal{B}_{\mathbb{C}^\times} := \left\{ \prod_{i \in I, a \in \mathbb{C}^\times} Y_{i,a}^{m_{i,a}} \mid m_{i,a} \geq 0 \right\} \subset \mathcal{Y}_{\mathbb{C}^\times}$ dominant monomials.

Classification of simple modules [Chari-Pressley]

There is a one-to-one correspondence :

$$\begin{array}{ccc} \{\text{simple modules in } \mathcal{C}\} / \sim & \leftrightarrow & \mathcal{B}_{\mathbb{C}^\times} \\ \Psi & & \Psi \\ [L(m)] & \leftrightarrow & m \end{array}$$

\exists an “affine analogue” $A_{i,a} \in \mathcal{Y}_{\mathbb{C}^\times}$ of e^{α_i} , α_i simple root.

Type $A_n^{(1)}$

$$A_{i,a} = Y_{i,aq^{-1}} Y_{i,aq} Y_{i-1,a}^{-1} Y_{i+1,a}^{-1} \quad (\Leftrightarrow e^{\alpha_i} = e^{2\varpi_i - \varpi_{i-1} - \varpi_{i+1}})$$

$$(Y_{0,a} = Y_{n+1,a} := 1, e^{\varpi_0} = e^{\varpi_{n+1}} := 1.)$$

q-characters (2)

Set $\mathcal{B}_{\mathbb{C}^\times} := \left\{ \prod_{i \in I, a \in \mathbb{C}^\times} Y_{i,a}^{m_{i,a}} \mid m_{i,a} \geq 0 \right\} \subset \mathcal{Y}_{\mathbb{C}^\times}$ dominant monomials.

Classification of simple modules [Chari-Pressley]

There is a one-to-one correspondence :

$$\begin{array}{ccc} \{\text{simple modules in } \mathcal{C}\} / \sim & \leftrightarrow & \mathcal{B}_{\mathbb{C}^\times} \\ \Psi & & \Psi \\ [L(m)] & \leftrightarrow & m \end{array}$$

\exists an “affine analogue” $A_{i,a} \in \mathcal{Y}_{\mathbb{C}^\times}$ of e^{α_i} , α_i simple root.

Type $B_n^{(1)}$

$$A_{i,a} = \begin{cases} Y_{i,aq^{-2}} Y_{i,aq^2} Y_{i-1,a}^{-1} Y_{i+1,a}^{-1} & \text{if } i \leq n-2 \\ Y_{n-1,aq^{-2}} Y_{n-1,aq^2} Y_{n-2,a}^{-1} Y_{n,aq^{-1}}^{-1} Y_{n,aq}^{-1} & \text{if } i = n-1 \\ Y_{n,aq^{-1}} Y_{n,aq} Y_{n-1,a}^{-1} & \text{if } i = n. \end{cases}$$

$(Y_{0,a} := 1)$

q -characters (2)

Set $\mathcal{B}_{\mathbb{C}^\times} := \left\{ \prod_{i \in I, a \in \mathbb{C}^\times} Y_{i,a}^{m_{i,a}} \mid m_{i,a} \geq 0 \right\} \subset \mathcal{Y}_{\mathbb{C}^\times}$ dominant monomials.

Classification of simple modules [Chari-Pressley]

There is a one-to-one correspondence :

$$\begin{array}{ccc} \{\text{simple modules in } \mathcal{C}\} / \sim & \leftrightarrow & \mathcal{B}_{\mathbb{C}^\times} \\ \Psi & & \Psi \\ [L(m)] & \leftrightarrow & m \end{array}$$

\exists an “affine analogue” $A_{i,a} \in \mathcal{Y}_{\mathbb{C}^\times}$ of e^{α_i} , α_i simple root.

Define the partial ordering on the set of Laurent monomials in $\mathcal{Y}_{\mathbb{C}^\times}$ as

$$m \geq m' \Leftrightarrow m^{-1}m' \text{ is a product of } A_{i,a}^{-1}\text{'s.}$$

Theorem (Frenkel-Mukhin)

$$\chi_q(L(m)) = m + (\text{sum of terms lower than } m), \quad \forall m \in \mathcal{B}_{\mathbb{C}^\times}.$$

q -characters (3)

\mathcal{C}_\bullet := the full subcategory of \mathcal{C} such that

object : V with $\chi_q(V) \in \mathbb{Z}[Y_{i,q^r}^{\pm 1} \mid i \in I, r \in \mathbb{Z}] =: \mathcal{Y}$.

Properties

- \mathcal{C}_\bullet is a (non-semisimple) abelian \otimes -subcategory.
- $\mathcal{C} = \bigotimes_{a \in \mathbb{C}^\times / q^{\mathbb{Z}}} (\mathcal{C}_\bullet)_a$ ($(\mathcal{C}_\bullet)_a$ is obtained from \mathcal{C}_\bullet by shift of the spectral parameter by a).

From now on, we always work in \mathcal{C}_\bullet , and write

$$Y_{i,r} := Y_{i,q^r} \quad A_{i,r} := A_{i,q^r} \quad \mathcal{B} := \mathcal{B}_{\mathbb{C}^\times} \cap \mathcal{Y}.$$

Example

- $\mathfrak{g} = \mathfrak{sl}_2, I = \{1\}, \chi_q(L(Y_{1,r})) = Y_{1,r} + Y_{1,r+2}^{-1} = Y_{1,r}(1 + A_{1,r+1}^{-1})$.
- $\mathfrak{g} = \mathfrak{so}_5, I = \{1, 2\},$
 $\chi_q(L(Y_{1,r})) = Y_{1,r} + Y_{2,r+1}Y_{2,r+3}Y_{1,r+4}^{-1} + Y_{2,r+1}Y_{2,r+5}^{-1} + Y_{1,r+2}Y_{2,r+3}^{-1}Y_{2,r+5}^{-1} + Y_{1,r+6}^{-1}$.

q -characters (3)

\mathcal{C}_\bullet := the full subcategory of \mathcal{C} such that

object : V with $\chi_q(V) \in \mathbb{Z}[Y_{i,q^r}^{\pm 1} \mid i \in I, r \in \mathbb{Z}] =: \mathcal{Y}$.

Properties

- \mathcal{C}_\bullet is a (non-semisimple) abelian \otimes -subcategory.
- $\mathcal{C} = \bigotimes_{a \in \mathbb{C}^\times / q^{\mathbb{Z}}} (\mathcal{C}_\bullet)_a$ ($(\mathcal{C}_\bullet)_a$ is obtained from \mathcal{C}_\bullet by shift of the spectral parameter by a).

From now on, we always work in \mathcal{C}_\bullet , and write

$Y_{i,r} := Y_{i,q^r}$ $A_{i,r} := A_{i,q^r}$ $\mathcal{B} := \mathcal{B}_{\mathbb{C}^\times} \cap \mathcal{Y}$.
For $m = \prod_{i \in I, r \in \mathbb{Z}} Y_{i,r}^{u_{i,r}} \in \mathcal{B}$, a **standard module** is defined as

$$M(m) := \bigotimes_{r \in \mathbb{Z}} \overrightarrow{\bigotimes}_{i \in I} L(Y_{i,r})^{\otimes u_{i,r}}.$$

$\rightsquigarrow \{[L(m)] \mid m \in \mathcal{B}\}$ and $\{[M(m)] \mid m \in \mathcal{B}\}$ are \mathbb{Z} -bases of $K(\mathcal{C}_\bullet)$.

Quantum Grothendieck rings (1)

We follow Hernandez's algebraic construction of quantum Grothendieck rings here.

Remark

\exists other (geometric) constructions given by Varagnolo-Vasserot and Nakajima for ADE cases, and all constructions produce equivalent rings in these cases.

First, we prepare a deformation \mathcal{Y}_t of the ambient Laurent polynomial ring \mathcal{Y} .

$\rightsquigarrow \mathcal{Y}_t$ is a $\mathbb{Z}[t^{\pm 1/2}]$ -algebra such that

- generators : $\tilde{Y}_{i,r}$ ($i \in I, r \in \mathbb{Z}$) and their inverses $\tilde{Y}_{i,r}^{-1}$
- relations : $\tilde{Y}_{i,r}$'s mutually **t -commute**.

e.g. $B_2^{(1)}$ -case : $\tilde{Y}_{1,r+2}\tilde{Y}_{1,r} = t\tilde{Y}_{1,r}\tilde{Y}_{1,r+2}$, $\tilde{Y}_{1,r+5}\tilde{Y}_{2,r} = t^{-1}\tilde{Y}_{2,r}\tilde{Y}_{1,r+5}, \dots$

Quantum Grothendieck rings (2)

There exists a \mathbb{Z} -algebra homomorphism $\text{ev}_{t=1}: \mathcal{Y}_t \rightarrow \mathcal{Y}$ given by

$$t^{1/2} \mapsto 1 \qquad \tilde{Y}_{i,r} \mapsto Y_{i,r}.$$

This map is called **the specialization** at $t = 1$.

There exists a \mathbb{Z} -algebra anti-involution $\overline{(\cdot)}$ on \mathcal{Y}_t given by

$$t^{1/2} \mapsto t^{-1/2} \qquad \tilde{Y}_{i,r} \mapsto t^{-1} \tilde{Y}_{i,r}.$$

This map is called **the bar-involution**.

$\forall m \in \mathcal{Y}$ monomial $\rightsquigarrow \exists! \underline{m} \in \mathcal{Y}_t$ monomial (with coefficient in $t^{\mathbb{Z}/2}$) such that $\overline{\underline{m}} = \underline{m}$. (e.g. $\underline{Y}_{i,r} = t^{-1/2} \tilde{Y}_{i,r}$.) Set $\tilde{A}_{i,r} := \underline{A_{i,r}}$.

Quantum Grothendieck rings (3)

For $i \in I$, set

$$K_{i,t} := \langle \tilde{Y}_{i,r}(1 + t\tilde{A}_{i,r+r_i}^{-1}), \tilde{Y}_{j,r}^{\pm 1} \mid j \in I \setminus \{i\}, r \in \mathbb{Z} \rangle_{\mathbb{Z}[t^{\pm 1/2}]\text{-alg.}} \subset \mathcal{Y}_t.$$

Define the quantum Grothendieck ring of \mathcal{C}_\bullet as

$$K_t(\mathcal{C}_\bullet) := \bigcap_{i \in I} K_{i,t}.$$

Remark

Indeed, $K_{i,t}$ = the kernel of a t -analogue of “the screening operator associated to $i \in I$ ” [Hernandez].

$\rightsquigarrow K_t(\mathcal{C}_\bullet)$ is an affine analogue of the space of “ W -invariant functions”.

Theorem (Varagnolo-Vasserot, Nakajima, Hernandez)

$$\text{ev}_{t=1}(K_t(\mathcal{C}_\bullet)) = \chi_q(K(\mathcal{C}_\bullet)).$$

(q, t) -characters (1)

\exists a $\mathbb{Z}[t^{\pm 1/2}]$ -basis $\{M_t(m) \mid m \in \mathcal{B}\}$ of $K_t(\mathcal{C}_\bullet)$ such that
 $\text{ev}_{t=1}(M_t(m)) = \chi_q(M(m))$ [Nakajima, Hernandez].
 $\rightsquigarrow M_t(m)$ is called **the (q, t) -character of $M(m)$** .

All $M_t(m)$ can be explicitly calculated once we know $M_t(Y_{i,0}), i \in I$.

Theorem (Nakajima (ADE cases), Hernandez (arbitrary))

$\exists! \{L_t(\underline{m}) \mid \underline{m} \in \mathcal{B}\}$ a $\mathbb{Z}[t^{\pm 1/2}]$ -basis of $K_t(\mathcal{C}_\bullet)$ such that

(S1) $\overline{L_t(\underline{m})} = L_t(\underline{m})$, and

(S2) $M_t(m) = L_t(m) + \sum_{m' < m} P_{m,m'}(t)L_t(m')$ with
 $P_{m,m'}(t) \in t^{-1}\mathbb{Z}[t^{-1}]$.

The element $L_t(m)$ is called **the (q, t) -character of $L(m)$** .

(q, t) -characters (2)

$$(S1) \overline{L_t(m)} = L_t(m) \quad (S2) M_t(m) = L_t(m) + \sum_{m' < m} P_{m, m'}(t) L_t(m'), \quad P_{m, m'}(t) \in t^{-1} \mathbb{Z}[t^{-1}]$$

Remark

The characterization properties (S1) and (S2) provide an inductive algorithm for computing $P_{m, m'}(t)$'s, called **Kazhdan-Lusztig algorithm**.

When \mathfrak{g} is of ADE type,

$$\text{ev}_{t=1}(L_t(m)) = \chi_q(L(m)) \text{ [Nakajima].}$$

Its proof is based on his geometric construction using quiver varieties, and it is valid only in ADE case. Moreover, in this case,

$$P_{m, m'}(t) \in t^{-1} \mathbb{Z}_{\geq 0}[t^{-1}] \text{ (positivity).}$$

(q, t) -characters (2)

(S1) $\overline{L_t(m)} = L_t(m)$ (S2) $M_t(m) = L_t(m) + \sum_{m' < m} P_{m,m'}(t)L_t(m')$, $P_{m,m'}(t) \in t^{-1}\mathbb{Z}[t^{-1}]$

Remark

The characterization properties (S1) and (S2) provide an inductive algorithm for computing $P_{m,m'}(t)$'s, called Kazhdan-Lusztig algorithm.

Conjecture (Hernandez)

For arbitrary cases, we also have

(1) $\forall m \in \mathcal{B}$, $\text{ev}_{t=1}(L_t(m)) = \chi_q(L(m))$. (2) $P_{m,m'}(t) \in t^{-1}\mathbb{Z}_{\geq 0}[t^{-1}]$.

If Conjecture (1) holds (in particular, in ADE cases), we have

$$[M(m)] = [L(m)] + \sum_{m' < m} P_{m,m'}(1)[L(m')] \text{ in } K(\mathcal{C}_\bullet).$$

Quantized coordinate algebra of type A_N

Let \mathcal{U}_v^- be the negative half of the QEA of type A_N over $\mathbb{Q}(v^{1/2})$.

(:= the $\mathbb{Q}(v^{1/2})$ -algebra with generators $\{f_i\}_{i=1,\dots,N}$, relations $\begin{cases} f_i^2 f_j - (v + v^{-1}) f_i f_j f_i + f_j f_i^2 = 0 & \text{if } |i - j| = 1 \\ f_i f_j - f_j f_i = 0 & \text{if } |i - j| > 1. \end{cases}$)

$\rightsquigarrow \mathcal{A}_v[N_-^{A_N}] \subset_{\mathbb{Z}[v^{\pm 1/2}]\text{-subalg}} \mathcal{U}_v^-$ the quantized coordinate algebra.

Property

$\mathbb{Q}(v^{\pm 1/2}) \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathcal{A}_v[N_-^{A_N}] \simeq \mathcal{U}_v^- \quad \mathbb{C} \otimes_{\mathbb{Z}[v^{\pm 1/2}]} \mathcal{A}_v[N_-^{A_N}] \simeq \mathbb{C}[N_-^{A_N}]$.

Here $N_-^{A_N} := \{(N + 1) \times (N + 1) \text{ unipotent lower triangular matrices}\}$.

- $\exists \text{ev}_{v=1}: \mathcal{A}_v[N_-^{A_N}] \rightarrow \mathbb{C}[N_-^{A_N}]$ a \mathbb{Z} -algebra homomorphism, called **the specialization** at $v = 1$.
- \exists an \mathbb{Z} -algebra anti-involution σ' on $\mathcal{A}_v[N_-^{A_N}]$, called **the (twisted) dual bar involution** (e.g. $v^{1/2} \mapsto v^{-1/2}$).

(:= the restriction of the \mathbb{Z} -algebra anti-involution on \mathcal{U}_v^- given by $v^{1/2} \mapsto v^{-1/2}, f_i \mapsto -f_i$.)

Dual canonical bases

Let $\mathbf{i} = (i_1, i_2, \dots, i_\ell)$ be a reduced word of the longest element w_0 of the Weyl group $W^{A_N} \simeq \mathfrak{S}_{N+1}$.

(e.g. if $N = 2$, then $\mathbf{i} = (1, 2, 1)$ or $(2, 1, 2)$.)

Dual canonical bases

Let $\mathbf{i} = (i_1, i_2, \dots, i_\ell)$ be a reduced word of the longest element w_0 of the Weyl group $W^{A_N} \simeq \mathfrak{S}_{N+1}$. Let Δ_+ be the set of positive roots of type A_N .

$\rightsquigarrow \exists \{ \widetilde{F}^{\text{up}}(\mathbf{c}, \mathbf{i}) \mid \mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Delta_+} \}$ a $\mathbb{Z}[v^{\pm 1/2}]$ -basis of $\mathcal{A}_v[N_-^{A_N}]$ depending on \mathbf{i} , which is an analogue of **the (dual) PBW-basis associated to \mathbf{i}** [Lusztig].

Theorem (Lusztig, Saito, Kimura)

- $\exists ! \widetilde{\mathbf{B}}^{\text{up}} := \{ \widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}) \mid \mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Delta_+} \}$ a $\mathbb{Z}[v^{\pm 1/2}]$ -basis of $\mathcal{A}_v[N_-^{A_N}]$ such that
 - (B1) $\sigma'(\widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i})) = \widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i})$, and
 - (B2) $\widetilde{F}^{\text{up}}(\mathbf{c}, \mathbf{i}) = \widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}) + \sum_{\mathbf{c}' < \mathbf{c}} p_{\mathbf{c}, \mathbf{c}'}(v) \widetilde{G}^{\text{up}}(\mathbf{c}', \mathbf{i})$ with $p_{\mathbf{c}, \mathbf{c}'}(v) \in v\mathbb{Z}[v]$.
- $\widetilde{\mathbf{B}}^{\text{up}}$ does not depend on the choice of \mathbf{i} .

The basis $\widetilde{\mathbf{B}}^{\text{up}}$ is called **the (normalized) dual canonical basis**.

Positivities

$$(B1) \sigma'(\widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i})) = \widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}) \quad (B2) \widetilde{F}^{\text{up}}(\mathbf{c}, \mathbf{i}) = \widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}) + \sum_{\mathbf{c}' \neq \mathbf{c}} p_{\mathbf{c}, \mathbf{c}'}(v) \widetilde{G}^{\text{up}}(\mathbf{c}', \mathbf{i}), p_{\mathbf{c}, \mathbf{c}'}(v) \in v\mathbb{Z}[v]$$

Theorem (Lusztig (i “adapted”), Kato, McNamara (arbitrary), (O. arbitrary))

$$p_{\mathbf{c}, \mathbf{c}'}(v) \in \mathbb{Z}_{\geq 0}[v].$$

Theorem (Lusztig)

For $\mathbf{c}_1, \mathbf{c}_2 \in \mathbb{Z}_{\geq 0}^{\Delta_+}$, write

$$\widetilde{G}^{\text{up}}(\mathbf{c}_1, \mathbf{i}) \widetilde{G}^{\text{up}}(\mathbf{c}_2, \mathbf{i}) = \sum_{\mathbf{c}} c_{\mathbf{c}_1, \mathbf{c}_2}^{\mathbf{c}} \widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}).$$

Then $c_{\mathbf{c}_1, \mathbf{c}_2}^{\mathbf{c}} \in \mathbb{Z}_{\geq 0}[v^{\pm 1/2}]$.


Hernandez-Leclerc isomorphisms in type $A_N^{(1)}$ (1)

Assume that $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ is of type $A_N^{(1)}$ ($I = \{1, \dots, N\}$).


Define $J_{Q', A_N^{(1)}}$ by

$$J_{Q', A_N^{(1)}} := \{(i, -i+1-2k) \in I \times \mathbb{Z} \mid k = 0, 1, \dots, 2n-i-1 \text{ and } i \in I\}.$$

$N = 4$



$N = 5$



Hernandez-Leclerc isomorphisms in type $A_N^{(1)}$ (1)

Assume that $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ is of type $A_N^{(1)}$ ($I = \{1, \dots, N\}$).

Define $J_{Q', A_N^{(1)}}$ by

$$J_{Q', A_N^{(1)}} := \{(i, -i+1-2k) \in I \times \mathbb{Z} \mid k = 0, 1, \dots, 2n-i-1 \text{ and } i \in I\}.$$

Set

$$\mathcal{B}_{Q', A_N^{(1)}} := \left\{ \prod_{(i,r)} Y_{i,r}^{u_{i,r}} \in \mathcal{B} \mid u_{i,r} \neq 0 \text{ only if } (i,r) \in J_{Q', A_N^{(1)}} \right\},$$

$\mathcal{C}_{Q', A_N^{(1)}}$:= the full subcategory of \mathcal{C}_\bullet such that

$$\underline{\text{object}} : V \text{ with } [V] \in \sum_{m \in \mathcal{B}_{Q', A_N^{(1)}}} \mathbb{Z}[L(m)].$$

Lemma (Hernandez-Leclerc)

$\mathcal{C}_{Q', A_N^{(1)}}$ is an abelian \otimes -subcategory.

Hernandez-Leclerc isomorphisms in type $A_N^{(1)}$ (2)

Set

$$K_t(\mathcal{C}_{\mathcal{Q}', A_N^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathcal{Q}', A_N^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathcal{Q}', A_N^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

Lemma

$K_t(\mathcal{C}_{\mathcal{Q}', A_N^{(1)}})$ is a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra of $K_t(\mathcal{C}_\bullet)$.

$\rightsquigarrow K_t(\mathcal{C}_{\mathcal{Q}', A_N^{(1)}})$ is called the quantum Grothendieck ring of $\mathcal{C}_{\mathcal{Q}', A_N^{(1)}}$.

Hernandez-Leclerc isomorphisms in type $A_N^{(1)}$ (2)

Set

$$K_t(\mathcal{C}_{\mathcal{Q}', A_N^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathcal{Q}', A_N^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathcal{Q}', A_N^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

Write

$$J_{\mathcal{Q}', A_N^{(1)}} = \{(i_s, r_s) \mid s = 1, \dots, \ell (= N(N+1)/2)\} \text{ with } r_1 \geq \dots \geq r_\ell.$$

$\rightsquigarrow \mathbf{i}_{\mathcal{Q}'} := (i_1, i_2, \dots, i_\ell)$ is a reduced word of $w_0 \in W^{A_N}$.

Remark

The reduced word $\mathbf{i}_{\mathcal{Q}'}$ depends on the choice of the total ordering on $J_{\mathcal{Q}', A_N^{(1)}}$. However, its “commutation class” is uniquely determined.

The following results does not depend on this choice.

This $\mathbf{i}_{\mathcal{Q}'}$ is “**adapted to \mathcal{Q}'** ”.

Hernandez-Leclerc isomorphisms in type $A_N^{(1)}$ (2)

$$K_t(\mathcal{C}_{\mathcal{Q}', A_N^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathcal{Q}', A_N^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathcal{Q}', A_N^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

$$J_{\mathcal{Q}', A_N^{(1)}} = \{(l_s, r_s) \mid s = 1, \dots, \ell (= N(N+1)/2)\} \text{ with } r_1 \geq \dots \geq r_\ell.$$

$\rightsquigarrow \mathbf{i}_{\mathcal{Q}'} := (i_1, i_2, \dots, i_\ell)$ is a reduced word of $w_0 \in W^{A_N}$.

Theorem (Hernandez-Leclerc)

There exists a \mathbb{Z} -algebra isomorphism

$$\Phi_A : \mathcal{A}_v[N_-^{A_N}] \xrightarrow{\sim} K_t(\mathcal{C}_{\mathcal{Q}', A_N^{(1)}})$$

given by

$$v^{\pm 1/2} \mapsto t^{\mp 1/2} \quad \widetilde{F}^{\text{up}}(\mathbf{c}, \mathbf{i}_{\mathcal{Q}'}) \mapsto M_t(m(\mathbf{c})) \quad \forall \mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Delta_+},$$

here $m(\mathbf{c}) = \prod_{k=1}^{\ell} Y_{i_k, r_k}^{\mathbf{c}(s_{i_1} \dots s_{i_{k-1}} \alpha_{i_k})}$. Moreover,


$$\Phi_A(\widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}_{\mathcal{Q}'})) = L_t(m(\mathbf{c})). \quad \forall \mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Delta_+}.$$

Our isomorphisms (1)

Assume that $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ is of type $B_n^{(1)}$ ($I = \{1, \dots, n\}$).

Let $\tilde{I} := \{1, \dots, 2n - 1\}$. Define $\tilde{J}_{\mathcal{Q}, B_n^{(1)}}$ by

$$\begin{aligned} \tilde{J}_{\mathcal{Q}, B_n^{(1)}} := & \{(i, -i + 2 - 2k) \mid k = 0, \dots, 2n - 1 - i \text{ and } i = n + 1, \dots, 2n - 1\} \\ & \cup \{(n, -n + \frac{3}{2} - k) \mid k = 0, \dots, 2n - 2\} \\ & \cup \{(i, -i + 1 - 2k) \mid k = 0, \dots, 2n - 2 - i \text{ and } i = 1, \dots, n - 1\}. \end{aligned}$$




Our isomorphisms (1)

Assume that $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ is of type $B_n^{(1)}$ ($I = \{1, \dots, n\}$).

Let $\tilde{I} := \{1, \dots, 2n - 1\}$. Define $\tilde{J}_{\mathcal{Q}, B_n^{(1)}}$ by

$$\begin{aligned} \tilde{J}_{\mathcal{Q}, B_n^{(1)}} := & \{(i, -i + 2 - 2k) \mid k = 0, \dots, 2n - 1 - i \text{ and } i = n + 1, \dots, 2n - 1\} \\ & \cup \{(n, -n + \frac{3}{2} - k) \mid k = 0, \dots, 2n - 2\} \\ & \cup \{(i, -i + 1 - 2k) \mid k = 0, \dots, 2n - 2 - i \text{ and } i = 1, \dots, n - 1\}. \end{aligned}$$

$n = 5$



Our isomorphisms (1)

Assume that $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ is of type $B_n^{(1)}$ ($I = \{1, \dots, n\}$).

Let $\tilde{I} := \{1, \dots, 2n - 1\}$. Define $\tilde{J}_{\mathcal{Q}, B_n^{(1)}}$.

Consider the map $\tilde{I} \rightarrow I, i \mapsto \bar{i} := \begin{cases} i & \text{if } i \leq n, \\ 2n - i & \text{if } i > n. \end{cases}$ “folding”

Set

$$\mathcal{B}_{\mathcal{Q}, B_n^{(1)}} := \left\{ \prod_{(i,r)} Y_{i,r}^{u_{i,r}} \in \mathcal{B} \mid \begin{array}{l} u_{i,r} \neq 0 \text{ only if } (i,r) = (\bar{i}, 2s) \\ \text{for some } (i,s) \in \tilde{J}_{\mathcal{Q}, B_n^{(1)}} \end{array} \right\},$$

$\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$:= the full subcategory of \mathcal{C}_\bullet such that

$$\underline{\text{object}} : V \text{ with } [V] \in \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[L(m)].$$

Lemma (Oh-Suh, Hernandez-O.)

$\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ is an abelian \otimes -subcategory.

Our isomorphisms (1)


Assume that $\mathcal{U}_q(\mathcal{L}\mathfrak{g})$ is of type $B_n^{(1)}$ ($I = \{1, \dots, n\}$).

Let $\tilde{I} := \{1, \dots, 2n - 1\}$.

Consider the map $\tilde{I} \rightarrow I, i \mapsto \bar{i} := \begin{cases} i & \text{if } i \leq n, \\ 2n - i & \text{if } i > n. \end{cases}$ “folding”

Set

$$\mathcal{B}_{\mathcal{Q}, B_n^{(1)}} := \left\{ \prod_{(i,r)} Y_{i,r}^{u_{i,r}} \in \mathcal{B} \mid \begin{array}{l} u_{i,r} \neq 0 \text{ only if } (i,r) = (\bar{i}, 2s) \\ \text{for some } (i,s) \in \tilde{J}_{\mathcal{Q}, B_n^{(1)}} \end{array} \right\}.$$



Our isomorphisms (2)

Set

$$K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

Lemma

$K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$ is a $\mathbb{Z}[t^{\pm 1/2}]$ -subalgebra of $K_t(\mathcal{C}_\bullet)$.

$\rightsquigarrow K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$ is called the quantum Grothendieck ring of $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$.

Our isomorphisms (2)

Set

$$K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

Write

$$\tilde{J}_{\mathcal{Q}, B_n^{(1)}} = \{(v_s, r_s) \mid s = 1, \dots, \ell(= 2n(2n-1)/2)\} \text{ with } r_1 \geq \dots \geq r_\ell.$$

$\rightsquigarrow \mathbf{i}_{\mathcal{Q}}^{\text{tw}} := (v_1, v_2, \dots, v_\ell)$ is a reduced word of $w_0 \in W^{A_{2n-1}}$.

Remark

The reduced word $\mathbf{i}_{\mathcal{Q}}^{\text{tw}}$ depends on the choice of the total ordering on $J_{\mathcal{Q}, B_n^{(1)}}$. However, its “commutation class” is uniquely determined. The following results does not depend on this choice. This $\mathbf{i}_{\mathcal{Q}}^{\text{tw}}$ is always “**non-adapted**”.

Our isomorphisms (2)


Set

$$K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathcal{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

$$\tilde{J}_{\mathcal{Q}, B_n^{(1)}} = \{(\iota_s, r_s) \mid s = 1, \dots, \ell (= 2n(2n-1)/2)\} \text{ with } r_1 \geq \dots \geq r_\ell.$$

$\rightsquigarrow \mathbf{i}_{\mathcal{Q}}^{\text{tw}} := (\iota_1, \iota_2, \dots, \iota_\ell)$ is a reduced word of $w_0 \in W^{A_{2n-1}}$.

In the following example, $\mathbf{i}_{\mathcal{Q}}^{\text{tw}} = (1, 2, 3, 1, 4, 3, 2, 5, 3, 1, 4, 3, 2, 3, 1)$ etc.



Our isomorphisms (2)

$$K_t(\mathcal{C}_{\mathbb{Q}, B_n^{(1)}}) := \sum_{m \in \mathcal{B}_{\mathbb{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] M_t(m) = \sum_{m \in \mathcal{B}_{\mathbb{Q}, B_n^{(1)}}} \mathbb{Z}[t^{\pm 1/2}] L_t(m).$$

$$\tilde{J}_{\mathbb{Q}, B_n^{(1)}} = \{(l_s, r_s) \mid s = 1, \dots, \ell (= 2n(2n-1)/2)\} \text{ with } r_1 \geq \dots \geq r_\ell.$$

$$\rightsquigarrow \mathbf{i}_{\mathbb{Q}}^{\text{tw}} := (i_1, i_2, \dots, i_\ell) \text{ is a reduced word of } w_0 \in W^{A_{2n-1}}.$$

Theorem (Hernandez-O.)

There exists a \mathbb{Z} -algebra isomorphism

$$\Phi_B : \mathcal{A}_v[N_-^{A_{2n-1}}] \xrightarrow{\sim} K_t(\mathcal{C}_{\mathbb{Q}, B_n^{(1)}})$$

given by

$$v^{\pm 1/2} \mapsto t^{\mp 1/2} \quad \widetilde{F}^{\text{up}}(\mathbf{c}, \mathbf{i}_{\mathbb{Q}}^{\text{tw}}) \mapsto M_t(m'(\mathbf{c})) \quad \forall \mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Delta_+},$$

here $m'(\mathbf{c}) = \prod_{k=1}^{\ell} Y_{i_k, r_k}^{\mathbf{c}(s_{i_1} \cdots s_{i_{k-1}} \alpha_{i_k})}$. Moreover,

$$\Phi_B(\widetilde{G}^{\text{up}}(\mathbf{c}, \mathbf{i}_{\mathbb{Q}}^{\text{tw}})) = L_t(m'(\mathbf{c})). \quad \forall \mathbf{c} \in \mathbb{Z}_{\geq 0}^{\Delta_+}.$$

Positivities in $\mathcal{C}_{\mathcal{Q}, \mathbb{B}_n^{(1)}}$

By our theorem, the positivities of the dual canonical bases $\widetilde{\mathbf{B}}^{\text{up}}$ can be transported to those of (q, t) -characters.

Corollary (Positivity of Kazhdan-Lusztig type polynomials)

For $m \in \mathcal{B}_{\mathcal{Q}, \mathbb{B}_n^{(1)}}$, write

$$M_t(m) = \sum_{m' \in \mathcal{B}_{\mathcal{Q}, \mathbb{B}_n^{(1)}}} P_{m, m'}(t) L_t(m').$$

as before. Then $P_{m, m'}(t) \in \mathbb{Z}_{\geq 0}[t^{-1}]$.

This is the affirmative answer to Conjecture (2) for $\mathcal{C}_{\mathcal{Q}, \mathbb{B}_n^{(1)}}$.

Corollary (Positivity of structure constants)

For $m_1, m_2 \in \mathcal{B}_{\mathcal{Q}, \mathbb{B}_n^{(1)}}$, write

$$L_t(m_1)L_t(m_2) = \sum_{m \in \mathcal{B}_{\mathcal{Q}, \mathbb{B}_n^{(1)}}} c_{m_1, m_2}^m L_t(m).$$

Then we have $c_{m_1, m_2}^m \in \mathbb{Z}_{\geq 0}[t^{\pm 1/2}]$.

Comparison with Kashiwara-Oh

The following remarkable theorem is recently proved by means of the celebrated *generalized quantum affine Schur-Weyl dualities*, which is developed by Kang, Kashiwara, Kim and Oh :

Theorem (Kashiwara-Oh '17)

There exists a \mathbb{Z} -algebra isomorphism

$$[\mathcal{F}]: \text{ev}_{v=1}(\mathcal{A}_v[N_-^{A_{2n-1}}]) \xrightarrow{\sim} K(\mathcal{C}_{\mathcal{Q}, \mathcal{B}_n^{(1)}})$$

which maps the dual canonical basis $\text{ev}_{v=1}(\tilde{\mathbf{B}}^{\text{up}})$ specialized at $v = 1$ to the set of classes of simple modules $\{[L(m)] \mid m \in \mathcal{B}_{\mathcal{Q}, \mathcal{B}_n^{(1)}}\}$.

Theorem (Hernandez-O.)

$$\Phi_{\mathcal{B}} \big|_{v=t=1} = [\mathcal{F}].$$

Comparison with Kashiwara-Oh

Theorem (Hernandez-O.)

$$\Phi_B |_{v=t=1} = [\mathcal{F}].$$

Remark

Our construction of Φ_B does not imply Kashiwara-Oh's theorem because, a priori,

- $\Phi_B |_{v=t=1}$ maps $\text{ev}_{v=1}(\tilde{\mathbf{B}}^{\text{up}})$ to $\{\text{ev}_{v=1}(L_t(m)) \mid m \in \mathcal{B}_{\mathcal{Q}, \mathbf{B}_n^{(1)}}\}$, but
- $[\mathcal{F}]$ maps $\text{ev}_{v=1}(\tilde{\mathbf{B}}^{\text{up}})$ to $\{[L(m)] \mid m \in \mathcal{B}_{\mathcal{Q}, \mathbf{B}_n^{(1)}}\}$,

(The coincidence of these images is nothing but Hernandez's conjecture (1) !) Hence our result and Kashiwara-Oh's result are independent.

Our comparison theorem above is proved by looking at the images of dual PBW-bases.

Comparison with Kashiwara-Oh

Theorem (Kashiwara-Oh '17)

There exists a \mathbb{Z} -algebra isomorphism

$$[\mathcal{F}]: \text{ev}_{v=1}(\mathcal{A}_v[N_-^{A_{2n-1}}]) \xrightarrow{\sim} K(\mathcal{C}_{\mathcal{Q}, \mathcal{B}_n^{(1)}})$$

which maps the dual canonical basis $\text{ev}_{v=1}(\tilde{\mathbf{B}}^{\text{up}})$ specialized at $v = 1$ to the set of classes of simple modules $\{[L(m)] \mid m \in \mathcal{B}_{\mathcal{Q}, \mathcal{B}_n^{(1)}}\}$.

Theorem (Hernandez-O.)

$$\Phi_{\mathcal{B}} \big|_{v=t=1} = [\mathcal{F}].$$

Corollary

$$\chi_{\mathcal{Q}}(L(m)) = \text{ev}_{t=1}(L_t(m)), \forall m \in \mathcal{B}_{\mathcal{Q}, \mathcal{B}_n^{(1)}}.$$

This is the affirmative answer to Conjecture (1) for $\mathcal{C}_{\mathcal{Q}, \mathcal{B}_n^{(1)}}$.

Comments on further results and proofs (1)

- There are several variants in the choices of the subcategories $\mathcal{C}_{\mathcal{Q}', A_N^{(1)}}$ and $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$. However the parallel results hold. (The choice in this talk is the case that \mathcal{Q}' and \mathcal{Q} are “equioriented”).
- By combining our Φ_B with Φ_A for $A_{2n-1}^{(1)}$, we can obtain a $\mathbb{Z}[v^{\pm 1/2}]$ -algebra isomorphism $K_t(\mathcal{C}_{\mathcal{Q}', A_{2n-1}^{(1)}}) \simeq K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$. For the choices of $\mathcal{C}_{\mathcal{Q}', A_{2n-1}^{(1)}}$ and $\mathcal{C}_{\mathcal{Q}, B_n^{(1)}}$ in this talk, we know **explicit correspondence of simple modules in terms of highest monomials.**

Key point

Highest monomial parametrization of simple modules =
PBW-parametrization of the dual canonical basis

Comments on further results and proofs (2)

Sketch of the proof of the existence of Φ_B

0) We have

- $K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$ “truncate” \hookrightarrow the quantum torus of *finitely many* variables.
- $\mathcal{A}_v[N_-^{A_{2n-1}}] \hookrightarrow$ the quantum torus arising from the “quantum initial seed” associated with $i_{\mathcal{Q}}^{\text{tw}}$ (\Leftarrow quantum cluster algebra).

- 1) Prove **the isomorphism between ambient tori** in Step 0. (Here we also use the cluster algebraic observation “ $A_{i,r}$ ’s are \hat{Y} -variables”)
- 2) Show **the coincidence between quantum T -system and quantum determinantal identities** (\Leftarrow mutation sequence. Every algebra generator appears as a cluster variable in this sequence).

Reference : arXiv:1803.06754v1

T -system

For $i \in I$, $r \in \mathbb{Z}$, $k \in \mathbb{Z}_{\geq 0}$, set $m_{k,r}^{(i)} := \prod_{s=1}^k Y_{i,r+2r_i(s-1)}$. ($m_{1,r}^{(i)} = Y_{i,r}$)

The quantum T -system of type B [Hernandez-O.]

$\exists \alpha, \beta \in \mathbb{Z}$ such that the following identity holds in $K_t(\mathcal{C}_{\mathcal{Q}, B_n^{(1)}})$:

$$L_t(m_{k,r}^{(i)})L_t(m_{k,r+2r_i}^{(i)}) = t^{\alpha/2}L_t(m_{k+1,r}^{(i)})L_t(m_{k-1,r+2r_i}^{(i)}) + t^{\beta/2}S_{k,r,t}^{(i)}.$$

Here, $S_{k,r,t}^{(i)} = \begin{cases} L_t(m_{k,r+2}^{(i-1)})L_t(m_{k,r+2}^{(i+1)}) & \text{if } i \leq n-2, \\ L_t(m_{k,r+2}^{(n-2)})L_t(m_{2k,r+1}^{(n)}) & \text{if } i = n-1, \\ L_t(m_{s,r+1}^{(n-1)})L_t(m_{s,r+3}^{(n-1)}) & \text{if } i = n \text{ and } k = 2s \text{ is even,} \\ L_t(m_{s+1,r+1}^{(n-1)})L_t(m_{s,r+3}^{(n-1)}) & \text{if } i = n \text{ and } k = 2s+1 \text{ is odd.} \end{cases}$ ($L_t(m_{*,*}^{(0)}) := 1$).

Example ($B_3^{(1)}$ -case)

- $L_t(m_{2,r}^{(1)})L_t(m_{2,r+4}^{(1)}) = tL_t(m_{3,r}^{(1)})L_t(m_{1,r+4}^{(1)}) + L_t(m_{2,r+2}^{(2)}).$
- $L_t(m_{3,r}^{(3)})L_t(m_{3,r+2}^{(3)}) = t^{1/2}L_t(m_{4,r}^{(3)})L_t(m_{2,r+2}^{(3)}) + t^{-1/2}L_t(m_{2,r+1}^{(2)})L_t(m_{1,r+3}^{(2)}).$